

Normal and Anomalous Heat Transport in One-Dimensional Classical Lattices

Tomaž Prosen¹ and David K. Campbell²

1. *Physics Department, Faculty of Mathematics and Physics, University of Ljubljana, Jadranska
19, 1111 Ljubljana, Slovenia*

2. *Departments of Physics and Electrical and Computer Engineering, College of Engineering,
Boston University, Boston, MA 02215, U.S.A*

(Published in: *Chaos* **15** (2005) 015117)

Abstract

We present analytic and numerical results on several models of one-dimensional (1D) classical lattices with the goal of determining the origins of anomalous heat transport and the conditions for normal transport in these systems. Some of the recent results in the literature are reviewed and several original new “toy” models are added that provide key elements to determine which dynamical properties are necessary and which are sufficient for certain types of heat transport. We demonstrate with numerical examples that chaos in the sense of positivity of Lyapunov exponents is neither necessary nor sufficient to guarantee normal transport in 1D lattices. Quite surprisingly, we find that in the absence of momentum conservation, even ergodicity of an isolated system is not necessary for the normal transport. Specifically, we demonstrate clearly the validity of the Fourier law in a pseudo-integrable particle chain.

PACS numbers: 44.10.+i, 05.45.-a, 05.60.+w, 05.70.Ln

I. LEAD PARAGRAPH

The study of how heat is transported through solids has a history dating back to antiquity and culminating, in the regime of classical physics, in the Fourier law of heat conduction, which asserts that the flux of heat (the “heat current”) is proportional to a constant times the gradient of the temperature. This “normal transport” of heat corresponds to “diffusion” of the heat through the system and is observed in most physical systems. In some mathematical models and in certain specially prepared experiments, one can observe “anomalous” transport of heat. Strictly speaking, one uses the term “anomalous” transport to describe anything that is not “normal,” up to and including the case in which the heat propagates ballistically through the system. Numerous prior studies have sought to understand the origins of both normal and anomalous transport and to isolate key features of the systems that lead to each type of transport. A particularly important set of studies developed from the original Fermi-Pasta-Ulam study of the (supposed!) equipartition of energy in a one-dimensional chain of coupled *nonlinear* classical oscillators. Just as the study of the time evolution of the FPU system revealed the surprising result that there was no apparent evolution towards equipartition but instead a tendency towards the recurrence of the initial state, so the study of transport in FPU-like systems revealed that their thermal conductivity was indeed anomalous. In the present article, we explore systematically the conditions that lead to normal or anomalous conductivity in classical, one-dimensional chains (“lattices”). We find an interesting interplay among the consequences of translation invariance (momentum conservation), nonlinearity, complete integrability, and deterministic chaos, and demonstrate the particularly surprising result that chaos is neither necessary nor sufficient for normal conductivity in these systems.

II. INTRODUCTION

The related issues of thermalization, transport, and heat conduction in one-dimensional (1D) classical lattices have been sources of continuing interest (and frustration!) for several generations of physicists, particularly since the pioneering computational study of Fermi, Pasta, and Ulam (FPU) revealed the remarkable “little discovery”¹ that even in a strongly nonlinear 1D lattice recurrences of the initial state prevented the equipartition of energy and consequent thermalization. The complex of questions that has developed from the FPU work involves the interrelations among equipartition of energy (the questions like: Is there equipartition? In which modes?), local thermal equilibrium (Does the system reach a well-defined temperature locally? If so, what is it?), and the transport of energy/heat (Does the system obey Fourier’s/Ficke’s heat law? If not, what is the nature of the abnormal transport?). Review articles spread over nearly three decades have provided snapshots of the understanding (and confusion) at different stages of this odyssey^{2–8}.

In this article, we will explore a subset of these questions dealing with the transport of energy/heat in 1D classical lattices. In particular, we will focus on the many studies that have attempted to verify the validity of Fourier’s law of heat conduction

$$\langle \vec{J} \rangle = -\kappa \nabla T \tag{1}$$

in 1D classical lattices or “chains.”

Here, κ is the transport coefficient of thermal conductivity and is supposed to be an *intensive* observable, *i.e.*, independent of the size of the system. Strictly speaking, κ is well defined only for a system that *obeys* Fourier’s law and for which a *linear* temperature gradient is established. Further, since in general κ is a function of temperature, the relative temperature variation across the chain should be small for κ to be truly constant.

In the literature, the dependence of $\kappa(L)$ on the size L of the system has also been used to characterize the (degree of) anomalous transport. However, the definition of κ for an anomalous conductor, where no *internal* temperature gradient may be established,

is highly ambiguous. Usually, one defines it as $\kappa_G(L) \equiv JL/\Delta T$ where ΔT is the total temperature drop between the two thermal baths. In this paper we shall call such a definition of κ_G a *global thermal conductivity*, in contrast to another definition of the so-called *local thermal conductivity* $\kappa_L = J/\nabla T$, which may depend on the position in the lattice, since $\nabla T = dT(x)/dx$ and is unique only in the case of a linear temperature profile. We note that the global and local thermal conductivities are proportional, namely $\kappa_L = c\kappa_G$ where c does not depend on the size L , *if and only if* the temperature profile is scaling, i.e. $T(x) = \tau(x/L)$ where $\tau(\zeta)$ again does not depend on L . We will confront these subtle distinctions at several points in the ensuing discussion.

To place our work in the larger context, it is useful to recall that the issues of equipartition/thermal equilibration and heat conduction/energy flow, although clearly related, can in fact be studied separately. For instance, although an integrable system will never reach a thermal equilibrium ensemble unless it is started in one, the concept of “soliton statistical mechanics” is not an oxymoron: assuming that the system is in a thermal equilibrium ensemble, one can study what other aspects of statistical mechanics remain valid and, in particular, what role solitons play in the response of the system to external perturbations. Similarly, the study of thermal transport (Fourier’s/Ficke’s heat law), is the search for a *non-equilibrium steady state* in which heat flows across the system, and the flow is typically analyzed *assuming* the Green-Kubo formalism of linear response¹⁰, in terms of correlation functions in the thermal equilibrium (grand canonical) state, independent of whether a particular system can actually reach this state. Indeed, systems exhibiting “anomalous conductivity” are precisely those in which this analysis does *not* lead to Fourier’s law.

We shall focus here on heat conduction but will attempt to make clear whenever our results impact (or depend on) the existence of equipartition and local thermal equilibrium.

Previous studies have led to bewildering array of partial results and conjectures:

- harmonic chains¹¹, showing $\kappa_G \sim L^1$, a result understood by the stability of linear Fourier modes, absence of mode-mode coupling, with the rigorous consequence that

no thermal gradient can be formed in the system.

- integrable models⁶, $\kappa_G \sim L^1$, a result understood by the presence of *nonlinear* stable modes (the solitons) and complete interability⁶;
- non-integrable models with smooth potentials : (1) the FPU models, leading eventually to claim that chaos was necessary and sufficient for normal conductivity ($\kappa_G \sim \kappa_L \sim L^0$), Ref.⁷, a claim that has been countered by strong numerical evidence for anomalous conductivity in FPU chains ($\kappa_G \sim \kappa_L \sim L^\alpha$, where $\alpha \approx 0.37$)^{12,13}; (2) the diatomic Toda chain, where initial results claiming $\kappa_G \sim L^0$, Ref.¹⁴ have recently been refuted by a study showing $\alpha \approx 0.4$, Ref.¹⁵; (3) the Frenkel-Kontorova model, which shows (at least for low temperatures) $\kappa_G \sim \kappa_L \sim L^0$, Ref.¹⁶;
- non-interable models with hard-core potentials (1) the “ding-a-ling” model¹⁷ and (2) the “ding-dong” model¹⁸, both showing convincingly that $\kappa_G \sim \kappa_L \sim L^0$. As one of the oldest studies of this type we should perhaps also mention Northcote and Potts model¹⁹.

This bewildering array of results has recently been greatly clarified in a series of independent but overlapping studies. The numerical studies of Hu, Li, and Zhao¹⁶ and of Hatano¹⁵ show that *overall momentum conservation* appears to a key factor in anomalous transport in 1D lattices. Lepri et al^{20,21} and Hatano¹⁵ have argued that the anomalous transport in momentum conserving systems can be understood in terms of low frequency, long-wavelength “hydrodynamic modes” that exist in typical momentum conserving systems and that *hydrodynamic arguments may explain the exponents observed in FPU^{20,21} and diatomic Toda¹⁵ lattices*. Later on, several other papers appeared which tried to formalize the statement that momentum conservation implies anomalous transport^{22–24}.

In the present paper, we review these recent results and extend them on several fronts, using numerical simulations of several different models. First, we present a counterexample showing that converse of the above statement is not true: namely, anomalous conductivity

does *not* imply that the model is momentum conserving. Second, we present convincing numerical data showing that chaos is neither necessary nor sufficient for normal conductivity, providing counterexamples to prior claims⁷. Finally, we present some (hopefully well-motivated) speculations on the necessary and sufficient conditions for *normal* conductivity in 1D lattice systems.

The class of models we study is described by the quite general 1D classical lattice Hamiltonian with single-particle and two-particle interactions

$$H \equiv \sum_n \left(\frac{p_n^2}{2m_n} + U_n^{\text{os}}(q_n) + V_{n+1/2}^{\text{ip}}(q_{n+1} - q_n) \right) \quad (2)$$

Here, the particles moving in 1D have “absolute” coordinates $x_n = na + q_n$, where a is the average lattice spacing and the relative coordinates q_n designate the particles’ displacements from their equilibrium (average) positions. We will either consider the particles to be on a ring with periodic boundary conditions (in this case $x_N \equiv x_0 + Na, q_N \equiv q_0$), or we will place the system between two thermal reservoirs (at possibly different temperatures) by coupling the edge particles 1 and N to canonical stochastic “heat” baths. To present our analytic results in the most general context, we will consider arbitrary “disorder”: the masses, m_n , can depend on lattice stite n , as can both the *on-site* potential U_n^{os} (which represents a “phonon-fixed lattice” interaction) and the *inter-particle* potential $V_{n+1/2}^{\text{ip}}$ (=“phonon-phonon” interaction).

We begin our detailed discussion in Section II with a survey of several different models (mostly previously studied) that fit into the general class described by the Hamiltonian in Eq. (2). The models are presented in a general sense of increasing “normality,” starting with (highly pathological) linear/harmonic chains, moving to (still pathological) integrable models, and finally treating (supposedly) “normal” chaotic models. Already from this survey we obtain two important results. First, we show that momentum conservation is not *necessary* for anomalous transport. The harmonic *optical* chain, which does not conserve momentum but does exhibit anomalous transport, provides one clear example of this, and an integrable model first introduced by Izergin and Korepin provies another. Second, by

reviewing recent results in the FPU chain, which is known to exhibit chaos but also clearly displays anomalous conductivity, we can conclude that chaos is not sufficient for normal conductivity, in contrast to previous claims⁷. In Section III, we discuss some of the recent theoretical work on the relation between the total momentum conservation and anomalous heat transport. We show that one can relate momentum conservation to the divergence of a particular form of the Kubo formula corresponding to a non-equilibrium system of spatially uniform density. However, uniform pressure, as required by the stationary state, produces the gradient of the density (or gradient of the chemical potential) which drives the ballistic current in the opposite direction to the temperature gradient. The two ballistic contributions thus cancel and the resulting next order term typically gives anomalous superdiffusive transport. In section IV we define and study the family of “bing-bang” models, which are constructed by considering purely *hard-wall* potentials U^{os} and V^{ip} . The bing-bang models are in fact equivalent to N -dimensional polyhedral billiards, which have vanishing Lyapunov exponents and therefore zero Kolmogorov-Sinai entropy. Hence, the bing-bang models are never chaotic in a sense of positive Lyapunov exponents. Nonetheless, we present convincing numerical evidence of *normal* (diffusive) heat transport in a (pseudo-integrable) non-chaotic but momentum non-conserving bing-bang model with an on-site potential. This demonstrates that *chaos*, at least in the sense of exponential instability of generic trajectories, is not necessary for the validity of the Fourier law. Taken together with the results from Section II, this shows the surprising result that chaos is neither necessary nor sufficient for normal conductivity in 1D classical lattices. In Section V we summarize our results and conclude by discussing possible precise necessary and sufficient conditions for normal and anomalous transport.

III. PARTIAL SURVEY OF PREVIOUS MODEL STUDIES

As this article is not a review of the very broad topic of anomalous transport^{8,25}, the ensuing survey of previous model studies is not intended to be exhaustive, but rather merely

illustrative, of the variety of models and results.

A. Linear Models

1. The linear acoustic chain

One of the first and few rigorous results on the issue of thermal conductivity is due to Rieder, Lebowitz and Lieb¹¹ who studied a linear (acoustic) chain of N (equal) harmonically coupled particles, where

$$U^{\text{os}}(q) = 0, \quad V^{\text{ip}}(q) = \frac{1}{2}\omega^2 q^2, \quad (3)$$

which are placed between two heat baths at temperatures T_1 and T_N . Rieder *et al.*¹¹ show that the transport is ballistic, with the average heat current J which is proportional to the *difference* of the temperatures of the two heat baths and independent of the size of the system

$$J = \gamma(T_N - T_1). \quad (4)$$

They also calculate the temperature profile T_j , $j = 1, 2 \dots N$, proving that it is asymptotically flat at the value of the average temperature $T(0 < j/N < 1) \rightarrow \frac{1}{2}(T_N + T_0)$, as $N \rightarrow \infty$. However, close to the boundaries of the system the temperature profile exhibits exponentially attenuated jumps to satisfy the boundary conditions. Surprisingly, these jumps are in the opposite direction from the temperatures of the corresponding heat baths (see Fig.1). Therefore, in the linear, *momentum-conserving* (acoustic) chain we find $\kappa_G = L^1$. But note that since the temperature profile is exponentially close to a flat profile, $\kappa_L \propto \exp(c_0 L)$.

2. The linear optical chain

Particularly in light of recent discussions on the role of on-site potential and total momentum conservation in heat conduction^{16,22}, it is natural to pose the following simple question:

what happens to the analysis of Rieder *et al* if a harmonic (quadratic) on-site potential is added, thereby breaking the momentum conservation of the model but retaining its linearity? Remarkably, we have been unable to find a previous reference in which this question was studied, so we present the results briefly here. The model is effectively a discretized massive Klein-Gordon system or linear *optical* chain, with

$$U^{\text{os}}(q) = \frac{1}{2}\Omega^2 q^2, \quad V^{\text{ip}}(q) = \frac{1}{2}\omega^2 q^2. \quad (5)$$

The formalism of Ref.¹¹ applies to the general class of linear chains, so it can be applied to the case of the linear on-site potential (5). Because the analysis is straightforward, we merely quote the results here. The momentum *non-conserving* linear optical chain also exhibits ballistic ($\kappa_G \sim L^1$) heat transport. Indeed, this optical model behaves in essentially the same way as the acoustic chain. The general shape of the temperature profile [eq.(4.2) of Ref.¹¹] and the expression of the heat current [eq.(4.6) of Ref.¹¹] remain exactly the same, the only difference being that the dimensionless parameter $\nu \propto \omega^2$ of the linear acoustic chain¹¹ is replaced by

$$\tilde{\nu} = \nu + \frac{\Omega^2}{\omega^2}, \quad (6)$$

for the linear optical chain. Note that this simple analysis already provides a non-trivial result: anomalous transport does *not* imply momentum conservation, so that momentum conservation is not a necessary condition for anomalous transport.

We should note that linear chains are also highly pathological in the additional sense that many of their dynamical properties depend on the spectral properties of model of the heat baths. This is a simple consequence of the existence of time conserved normal modes. It has been pointed out in Ref.²⁶, that in the case of the linear chain with disordered masses, one can obtain super diffusive $\alpha > 0$, diffusive $\alpha = 0$, or even sub-diffusive $\alpha < 0$, behavior for different choices of heat baths.

B. Nonlinear but Integrable Models

1. The Toda Lattice

As an example of a nonlinear but integrable momentum-conserving model, we consider the celebrated Toda lattice^{27,2}, for which

$$U^{\text{os}}(q) = 0, \quad V^{\text{ip}}(q) = \exp(-q). \quad (7)$$

Mokross and Büttner²⁸ have shown by numerical simulation that the temperature profile is almost flat and that the heat current is proportional to the difference of the bath temperatures (4). Hence they find numerical evidence for ballistic heat transport, with $\kappa_G(L) \propto L^1$, and $\kappa_L(L) \propto \exp(c_0 L)$, for this *integrable* and *momentum-conserving* but *non-linear* chain. For a discussion of the role of solitons in this anomalous transport, see Ref.⁶, and for the propagation of shock waves, see Ref.²⁹.

2. The Izergin-Korepin discrete sine-Gordon model

The natural class of models to consider next is a non-linear but integrable chain that does not conserve momentum, *i.e.*, has $U^{\text{os}}(q) \neq 0$. There are not many known models with such properties. However, one such model has been proposed by Izergin and Korepin³⁰ in the context of ultraviolet regularization of integrable quantum field theories in 1+1 dimensions. They introduced a spatially discrete version of the famous $\sin(\hbar)$ -Gordon model (we shall call this IK-SG model) which has the following properties: it preserves the integrability of the continuum limit and it has an on-site potential ($\sin \phi$ (for unimodular complex field) or $\sinh \phi$ (for real field)). The classical Hamiltonian formulation of the model has been described in detail by Tarasov in Ref.³¹. For simplicity, we consider here the case of real field variables $\phi_j^\pm(t)$, $j \in \mathbb{Z}$, which corresponds in the continuum limit to the \sinh -Gordon model with a confining anharmonic on-site potential. Unfortunately, the model cannot be cast exactly in the usual kinetic plus potential energy form of (2) except as an approximation to leading

order in lattice discreteness (see below). Nonetheless, the model provides important insight into the question of the role of U^{os} in anomalous conductivity. Referring to Ref.³¹, we find that the Hamiltonian of the discrete IK-SG model reads

$$H = \sum_n h_n, \quad h_n = h_n^+ + h_n^-, \quad (8)$$

$$h_n^\pm = \frac{(\phi_{n\mp 1}^+ \cos \delta - \phi_{n\pm 1}^- \sin \delta)(\phi_n^- \cos \delta - \phi_n^+ \sin \delta)}{(1 + \phi_{n\pm 1}^- \phi_n^+)(1 + \phi_{n\mp 1}^+ \phi_n^-)} + \frac{2}{\sin 2\delta}$$

where the parameter δ is related to a *lattice spacing* Δ via $\Delta = 2 \sin 2\delta$. For the *real* field variables ϕ_n^\pm , the Poisson bracket is defined as

$$\{\phi_n^-, \phi_m^+\} = \frac{1}{4 \cos 2\delta} \phi_n^+ \phi_n^- (\phi_n^+ \cos \delta - \phi_n^- \sin \delta)^2 \delta_{nm},$$

$$\{\phi_n^+, \phi_m^+\} = \{\phi_n^-, \phi_m^-\} = 0. \quad (9)$$

The field variables can be transformed to conventional (discretized) canonical fields, $u(x)$, $u_n = u(x_n)$ and canonical momentum field $\pi(x)$, $\pi_n = \pi(x_n)$, $p_n = \pi_n \Delta$, and $x_n = n\Delta$, by the transformation

$$\phi_n^\pm = \frac{\exp(\mp \frac{1}{2} u_n) \cos \delta + \exp(\pm \frac{1}{2} u_n) \sin \delta}{\sqrt{1 + \cosh(u_n) \sin 2\delta}} \exp(\frac{1}{4} p_n), \quad \{u_n, p_m\} = \delta_{nm} \quad (10)$$

In fact, the discrete Hamiltonian (8) can be written explicitly as an expansion in Δ of the continuum sinh-Gordon Hamiltonian as

$$H = \sum_n \left[\left(\frac{1}{2} \pi_n^2 + \frac{1}{8} (u_{n+1} - u_{n-1})^2 + \cosh u_n \right) \Delta + \frac{1}{2} \cosh(u_n) \pi_n \Delta^2 + \mathcal{O}(\Delta^3) \right]. \quad (11)$$

Izergin and Korepin have shown that the above discretized sinh-Gordon model (8) is completely solvable by the method of inverse scattering, and hence they have explicitly demonstrated its integrability.

We have used the above exact completely integrable model (8) in order to check the nature of energy transport in an IK-SG lattice. Since we have not found any natural intuitive way of coupling the variables ϕ_n^\pm (for $n = 1, N$) to the heat baths, we have decided to study the propagation of an initially localized pulse in an autonomous (isolated) lattice of very

large size N and with periodic boundary conditions $\phi_0^\pm \equiv \phi_N^\pm$. We start by initially exciting field variables at only one site (say $n = n_0 = N/2$) and setting all the others to the ground state (vacuum) value $\phi_{n \neq n_0}^\pm = 0$. Then we measure, numerically, after time t , the spatial spreading of the disturbance

$$\sigma^2(t) = \frac{1}{E} \sum_n (n - n_0)^2 h_n(t), \quad E = \sum_n h_n(t) = \text{const.} \quad (12)$$

For the numerical integration we use the adaptive step-size 4th order Runge-Kutta algorithm from Ref.³². Clearly, for *diffusive* energy transport, one would find $\sigma(t) \propto t^{1/2}$, whereas for *ballistic* transport, $\sigma(t) \propto t$. In figure 2 we show results of two numerical experiments with different values of the lattice parameter (δ or Δ); both show very clearly that (independent of the lattice parameter) the transport is ballistic, $\sigma(t) \propto t$.

This strongly suggests that integrability alone (the existence of an infinite number of independent conserved quantities in one-to-one correspondence with the set of degrees of freedom) is enough to yield anomalous (in fact, ballistic) heat transport, irrespective of the presence of on-site potential. This provides a second illustration, this time in a nonlinear context, of the result that momentum conservation is not necessary for anomalous transport.

C. Chaotic Models with smooth potentials

1. Chaotic but momentum conserving: the FPU models

In a series of recent studies^{12,13,20,21} of the celebrated FPU β chain^{1,7}, for which

$$U^{\text{os}}(q) = 0, \quad V^{\text{ip}}(q) = \frac{1}{2}q^2 + \frac{1}{4}\beta q^4, \quad (13)$$

several groups have shown by careful numerical analyses that despite strongly chaotic behavior (characterized by almost everywhere positive Lyapunov exponents), the model exhibits power-law divergence of thermal conductivity in thermodynamic limit, $\kappa_L \sim L^\alpha$, with $\alpha \approx 0.4$. Later, Hatano has shown¹⁵ that for the diatomic Toda chain (same as (7) but for different (dimerized) masses, $m_{2n} = m_2 \neq m_{2n+1} = m_1$) there exists practically the same

scaling with $\alpha \approx 0.4$ as for FPU models. More recently, a similar scaling has been shown for two other models: namely, for 1D chains where $V^{\text{ip}}(q)$ has the form of Lennard-Jones or Morse potentials³³. These results suggests possible universality of the scaling exponent $\alpha \approx 0.4$ (perhaps $2/5$) of the divergence of thermal conductivity for strongly chaotic, momentum conserving lattices. This universality has been partially explained

by Hatano¹⁵ using hydrodynamic arguments or by Lepri *et al*^{20,21,8} using mode-coupling theory⁴⁸.

However, recently Narayan and Ramaswamy²³ proposed another ‘thermodynamic’ approach to mode-mode coupling theory, which using a mapping to Burgers equation and the renormalization group predicts the universal exponent for the divergence of heat conductivity to be $\alpha = 1/3$. It should be mentioned that there exist other models with no-on-site potential which should conform to 1D hydrodynamics in thermodynamics, for example diatomic hard-point gas in 1D, which seems to have a smaller exponent α . Depending on the range and type of simulation^{34,24,35} one finds values in the range $\alpha \in [0.25, 0.35]$.

On the other hand, there exist certain momentum conserving particle chains which seem to exhibit normal heat conduction to a very high numerical accuracy. This is the case for the so-called rotator model^{36,37}, with no on-site potential $U^{\text{os}} \equiv 0$, and cosine interparticle potential $V^{\text{ip}}(q) = V_0 \cos q$. This model has a characteristic feature, namely it cannot support a non-vanishing pressure, and thus the infinite-wavelength phonons cannot carry any energy current²², but it is not clear why it should be excluded from the universality arguments²³. Thus there remain still unsettled issues in connection to the heat transport in momentum conserving though chaotic lattices. For some recent progress see e.g. Refs^{38,39}.

In any case, note that the existence of anomalous conductivity in models that exhibit strong chaos proves that chaos is not sufficient for normal conductivity, refuting earlier claims⁷. In Section IV, we shall show the still more surprising result that chaos is also not necessary for normal conductivity.

2. Chaotic but momentum nonconserving: the Frenkel-Kontorova model

Apart from a few more or less artificial models, such as ding-a-ling¹⁷ and ding-dong¹⁸ chains, which combine smooth and hard-core potentials, only one chaotic momentum non-conserving model with a smooth potential has been studied recently: namely, the Frenkel-Kontorova chain, with

$$U^{\text{os}}(q) = U_0 \cos q, \quad V^{\text{ip}}(q) = \frac{1}{2}q^2. \quad (14)$$

The numerical results of Hu, Li, and Zhao¹⁶ establish convincingly that this model exhibits *normal* heat transport— $\kappa_{\text{L}} \sim L^0$ —*provided* that the temperature is low enough: specifically, the thermal energy ($\sim k_B T$) must be smaller than the amplitude of the on-site potential barrier (U_0 in Eq. (14)). However, for temperatures well above U_0/k_B , the mean free path of the phonons becomes (exponentially) large, so the Fourier law is no longer simple to observe numerically. Very recently, Ref.⁴⁰ reports on extensive numerical simulations which appears to confirm the normal heat transport in FK chain even for temperatures above U_0/k_B .

IV. MOMENTUM CONSERVATION AND ANOMALOUS TRANSPORT

In this section we discuss some interesting and important connections between momentum conserving lattices and anomalous heat transport in 1D which follow from considering Kubo formula and the linear response theory.

In the absence of an on-site potential, the general Hamiltonian of Eq. (2) becomes

$$H_{\text{MC}} = \sum_{n=0}^{N-1} \left(\frac{1}{2m_n} p_n^2 + V_{n+1/2}(q_{n+1} - q_n) \right), \quad (15)$$

where $V_{n+1/2}(q)$ is an arbitrary (generally non-linear) interparticle interaction and the subscript “MC” stands for momentum conserving. As in Eq. (2) the inter-particle potential, $V_{n+1/2}(q_{n+1} - q_n)$, and the masses, m_n , may depend on the site label n . However there is now *no* “on-site” potential, $U^{\text{os}}(q_n)$, depending on the individual coordinates. Thus, H_{MC} is invariant under translations $q_n \rightarrow q_n + b$ for *arbitrary* b , and this spatial translational

symmetry corresponds to total momentum conservation. As before, we consider the (finite) system to be defined on a system of length $L = Na$ with periodic boundary conditions $(q_N, p_N) \equiv (q_0, p_0)$, where the actual particle positions are $x_n = na + q_n$.

We may write the Hamiltonian in Eqn.(15) as $H_{\text{MC}} = \sum_{n=0}^{N-1} h_{n+1/2}$, where $h_{n+1/2}$ is the Hamiltonian density

$$h_{n+1/2} = \frac{p_{n+1}^2}{4m_{n+1}} + \frac{p_n^2}{4m_n} + V_{n+1/2}(q_{n+1} - q_n), \quad (16)$$

Our aim is to estimate κ , the coefficient of thermal conductivity. Using the standard Kubo formula expression for κ of linear response (see e.g. Ref.⁴¹)

$$\kappa = \lim_{T \rightarrow \infty} \lim_{L \rightarrow \infty} \frac{\beta}{L} \int_{-T}^T dt \langle J(t)J \rangle_{\beta}, \quad (17)$$

then one can show, from very elementary arguments²², that $\kappa = \infty$ provided the thermodynamic pressure ϕ (i.e. average force between an arbitrary pair of adjacent particles) is non-vanishing. Here $\langle \rangle_{\beta}$ denotes a canonical phase space average at inverse temperature β . The approach of Ref.²² has been criticized^{25,23} by the claim that the heat current of the Kubo formula (17) needs a modification for the case of a system with Galilean invariance: Namely, the center of mass motion should be subtracted from the current, or one should put oneself into the frame where the total momentum is zero.

However, the formulation of Ref.²² was based on a slightly non-standard, though thermodynamically completely equivalent heat current, namely the so-called *lattice current*

$$J = \sum_{n=0}^{N-1} j_n \quad (18)$$

where j_n is the heat current density¹⁶, given by

$$j_n = \{h_{n+1/2}, h_{n-1/2}\} = \quad (19)$$

$$= \frac{p_n}{2m_n} (V'_{n+1/2}(q_{n+1} - q_n) + V'_{n-1/2}(q_n - q_{n-1})). \quad (20)$$

$\{, \}$ is the usual canonical Poisson bracket. Note that this form of j_n does correspond to the intuitive definition that the time rate of change of the energy at location n which should be

given by the (net) force $\sim (V'_{n+1/2}(q_{n+1} - q_n) + V'_{n-1/2}(q_n - q_{n-1}))$ times the velocity $\frac{p_n}{2m_n}$. More importantly, the current density j_n by construction satisfies the continuity equation

$$\frac{d}{dt}h_{n+1/2} = j_{n+1} - j_n. \quad (21)$$

Since the usual derivation of linear response and Kubo formula are based on the *real-space heat current* $J_{\text{rs}} = \sum_n \frac{1}{2}m_n v_n^3$, where $v_n = p_n/m_n$ is the velocity of n -th particle, one needs to rederive carefully the linear response formalism for the lattice current. This has been done in Ref.²⁴. Now, we wish to stress that the lattice current J and the real-space current J_{rs} are only equivalent if the center of mass motion is zero, precisely as required by the arguments of^{25,23}. In other words, adding a nonvanishing center-of-mass motion to the current J_{rs} , does not change the lattice current J . Still, as shown in Ref.²⁴ the Kubo formula (17) does not refer to the proper perturbation of the equilibrium state so it does not describe the steady heat current between two heat baths.

This result is quite interesting and consistent with the interpretation given in Ref.^{25,23}. The unmodified Kubo formula (17), with the current (19), is the correct one if one considers a particular kind of non-equilibrium initial state, namely such with a *spatially uniform density* (or uniform chemical potential). This we call the *isochoric* initial state, and in such a case we find rigorously²² that

$$\frac{\beta}{L} \int_0^T dt \langle J(t)J \rangle_\beta \geq \frac{t_0}{a\bar{m}} \phi^2 T, \quad (22)$$

so the transport is ballistic.

However, such a state cannot approximate a *steady state* of a large piece of the lattice between two heat baths at slightly different temperatures. In such a physical situation, a steady state is formed where the *pressure* is constant along the system and not the *density*. Typically, in colder regions of the lattice, the density is larger and vice versa. This we call the case of *isobaric* initial state. Then, due to non-uniform density, a gradient of chemical potential is established as well, which drives the heat current in the opposite direction. To leading order in L , the contributions to the heat current due to the temperature and chemical

potential gradients cancel each other. The difference of the two currents still produces the anomalous transport, since it decreases as $L^{\alpha-1}$. This has been demonstrated in a numerical experiment with high accuracy²⁴. The modified Kubo formula referring to the *isobaric* situation²⁴ reads

$$\kappa = \lim_{T \rightarrow \infty} \lim_{L \rightarrow \infty} \frac{\beta}{L} \int_{-T}^T dt \{ \langle J(t)J \rangle_{\beta} + \phi \langle V(t)J \rangle_{\beta} \}, \quad (23)$$

where $V = \sum_n v_n$ is the sum of all particles' velocities v_n .

Although we have learned that the issue of anomalous transport cannot be resolved easily solely only on the basis of momentum conservation, it is clear from this analysis that the momentum conservation is a key attribute of the problem. It is also interesting to observe that in the case of vanishing pressure $\phi = 0$ the isochoric and isobaric situations are described by identical Kubo formulae (17,23).

V. CHAOS AND NORMAL TRANSPORT: THE “BING-BANG” MODEL

The relationship between “chaos” and normal transport has been the subject of considerable interest, and it has even been claimed⁷ that chaos is both necessary and sufficient for the existence of normal transport. We have already shown that there exist models that exhibit both chaotic behavior and anomalous transport, so chaos is clearly not sufficient for normal transport. In this section, we establish the perhaps more surprising result that chaos is also not *necessary* for normal transport. We demonstrate this result using numerical studies of a 1D lattice model which, while exhibiting no chaos at all in the strict mathematical sense, nonetheless exhibits normal conductivity.

We call the particular class of many-body 1D classical lattices we study in this section the “*bing-bang*” models. As we shall see, they are manifestly non-chaotic in the sense of having Lyapunov exponents that are almost everywhere vanishing, but they are also non-integrable. The bing-bang models have infinite hard wall forms for both the *interparticle* potentials $V^{\text{ip}}(q)$, and the on-site potentials $U^{\text{os}}(q)$. If we fix the units such that the average

lattice spacing is equal to unity, $a = 1$, then the general (homogenous) bing-bang model would depend, apart from the masses m_n , on a triple of parameters $b, c, d \in \mathbb{R}^+$ which determine the potentials via

$$\begin{aligned} V^{\text{ip}}(q) &= \begin{cases} 0, & -1 \leq q \leq b; \\ \infty, & \text{otherwise} \end{cases} \\ U^{\text{os}}(q) &= \begin{cases} 0, & -c \leq q \leq d; \\ \infty, & \text{otherwise} \end{cases}. \end{aligned} \quad (24)$$

In fact, for a lattice of N particles, the bing-bang model can be identified with the motion of a point billiard particle inside an N -dimensional polytope. Since the boundary consists solely of $(N - 1)$ -dim. flat hyperplanes — *i.e.*, $-1 \leq q_n \leq b, -c \leq q_{n+1} - q_n \leq d$ — (almost) any orbit is marginally stable, *i.e.*, *parabolic* with zero asymptotic Lyapunov exponents. However, there is an infinite set (of vanishing Lebesgue measure in the full phase space) of very unstable orbits (which may also be called “diffractive” since they would produce diffraction of quantum mechanical waves) that eventually hit the (hyper)corners, edges, etc, either in finite future or in finite past. The simplest singular (infinitely unstable) orbits of this type are those that contain: (i) three particle collisions, or (ii) simultaneous collisions of a pair of particles and one of the walls.

Our main goal in studying such physically pathological systems is to gain insight into the extent to which the properties of exponential instability and metric chaos (which our bing-bang models do not possess) are necessary to produce normal transport (or perhaps anomalous transport with a “universal” exponent α if the total momentum is conserved). The insights gained in previous studies of models with pathological, hard wall potentials—such as the “ding-a-ling”¹⁷ and “ding-dong”¹⁸—provide ample motivation and justification for the current study. On the other hand, our results on decay of correlations (and, hence, perhaps on the mixing property) of non-hyperbolic (non-chaotic) systems with many-degrees of freedom should stimulate development of new tools in ergodic theory to deal with such systems in a rigorous way.

With respect to the global dynamical properties we define two subclasses of bing-bang

models.

(i) **Pseudo-integrable bing-bang (PIBB)** models in which the masses of all particles are equal (set to unity) $m_n = 1$. The pseudo-integrability of the model is easy to understand as follows. Upon interaction with interparticle potential $V(q_{n+1} - q_n)$ (interparticle collisions) the particles n and $n + 1$ just exchange their momenta/velocities, and upon interaction with the on-site potential $U(q_n)$ (collisions with a static hard wall) the momentum/velocity of the particle n just changes sign. Therefore, the dynamics in momentum space acts as a discrete group $(\mathcal{C}_2)^N \times \mathcal{S}_N$, and any symmetric function of the squares of momenta $f(p_1^2, \dots, p_N^2)$ is an invariant of motion. In fact, N independent analytic invariants of motion I_k can be systematically evaluated; they are the symmetric homogeneous polynomials of $\{p_n^2\}$

$$\begin{aligned}
I_1 &= \sum_{n=1}^L p_n^2 \\
I_2 &= \sum_{1 \leq n < m \leq L} p_n^2 p_m^2 \\
&\vdots \\
I_k &= \sum_{1 \leq n_1 < \dots < n_k \leq L} \prod_{l=1}^k p_{n_l}^2
\end{aligned} \tag{25}$$

Therefore, any orbit of a PIBB model lies on an invariant surface of dimension N in $2N$ -dimensional phase space, and the system is *not ergodic* on the entire energy surface. However, the invariant surface is not a simple N -torus as in the case of a completely integrable system, but is an object of (vastly) increasingly complex topology as N increases.

This is precisely the defining property of pseudo-integrable systems: namely, that there should exist a sufficient number of independent conservation laws for integrability but that the topology of invariant surfaces is more complicated than that of N -dimensional tori. This can only happen in the cases with singularities in the system: for instance, planar billiards in the shape of a polygon with angles that are rational multiples of π are the most commonly studied class of (two-dimensional) pseudo-integrable systems⁴².

In the case of our PIBB models, the dimensionality of the phase space $2N$ can be arbitrarily large and increasing N brings in new aspects and questions on dynamics that to the

best of our awareness have not yet been considered in the literature.

Our analysis below suggests that the topology of invariant surfaces can become arbitrarily complex as one approaches the thermodynamic limit $N \rightarrow \infty$ and that then such invariant surfaces can more and more densely and uniformly cover the entire energy surface, so that at the end, statistical mechanics cannot distinguish the system from a truly ergodic (and mixing) one. We suggest (but cannot prove) that the topological genus of the invariant surfaces in our PIBB lattices increases faster than any power of N . Since it is not essential to our current discussion, we shall leave the rigorous characterization of topology of invariant surfaces of high-dimensional PIBB models as a very interesting open mathematical problem.

(ii) **Ergodic bing-bang (EBB)** models. We have good heuristic arguments (and strong numerical evidence) to support the conjecture that the generic bing-bang model with different masses m_n (which should be generic, perhaps satisfying some irrationality conditions) is *ergodic*, since it corresponds to the motion inside N -dimensional polyhedral billiard. For example, it is rigorously known in the mathematical literature (see e.g.⁴² for a recent review, and references therein) that the set of ergodic polygonal billiards ($N = 2$) is (at least) a dense set in the set of all polygonal billiards with a fixed number of vertices. It seems fairly obvious that the same result should apply (even more likely) in higher dimensions $N > 2$. An even stronger result has been suggested recently:, namely that the generic polygonal billiard in the plane (e.g. triangular billiard with all angles having *irrational* ratio with π) is truly *mixing* dynamical system and therefore exhibits fast decay to statistical equilibrium from (almost) any initial state^{43,44}.

With respect to the key issue of total momentum conservation, we will study both *momentum conserving bing-bang models* (MC-PIBB or MC-EBB) which are characterized by $c, d = \infty$ (or saying that $U^{\text{os}} = 0$); and *momentum non-conserving bing-bang models* (MNC-PIBB or MNC-EBB), for which translational invariance is broken and generally at least one of parameters c, d is finite.

A. Momentum conserving bing-bang models

In this subsection we study two variants of momentum-conserving bing-bang models, one pseudo-integrable (MC-PIBB) and one ergodic (MC-EBB). Although they both behave as anomalous heat conductors we study both in order to show the subtle but important differences that are a consequence of qualitatively different positions in the *ergodic hierarchy* (pseudo-integrability versus ergodicity).

In our numerical simulations we will study either (i) an open bing-bang lattice between two thermal baths at temperatures T_L and T_R , or (ii) a closed bing-bang lattice with periodic boundary conditions where $N + 1 \equiv 1$. The thermal bath is realized as a wall (see e.g.⁴⁵) which works in the following way: when the edge particle (with label 1 or N) hits the reservoir, it collides inelastically so that the new momentum (being independent from the old one) is given with the probability distribution

$$d\mathcal{P}/dp_{1,N} \propto p_{1,N} \exp(-p_{1,N}^2/(2m_{1,N}T_{L,R})). \quad (26)$$

The velocity (momentum) prefactor takes into account the fact that the faster particles collide with the wall more often than the slow ones, so that the resulting velocity (momentum) distribution of the near-bath particles is indeed canonical Maxwellian (Gaussian).

Typical trajectories of the individual particles are shown in Fig.3 for the MC-PIBB model and in Fig. 4 for MC-EBB model, in order to give a reader an impression about the complex geometry of the orbits in bing-bang models despite their manifest non-chaoticity. In both cases, MC-PIBB and MC-EBB, the models are put between two heat reservoirs at the same temperature $T_L = T_R = T = 1$. In the following we will present results of numerical simulations of heat conduction in a non-equilibrium stationary state with the bath temperatures $T_L = 1$ and $T_R = 2$. We measure the average heat flux $J(L)$ through the system as the function of the size $L(= N, \text{ since } a = 1)$ and the kinetic temperature profile

$$T_n = \frac{\langle p_n^2 \rangle}{2m_n}.$$

First, we observe that temperature profile is typically a non-linear function (non-constant ∇T), although it has a scaling property: namely, the local temperature for a given system is only a function of the scaled coordinate n/N and reservoir temperatures

$$T_n = \tau(n/N, T_L, T_R). \quad (27)$$

In Figs.5, and 6, we show the sets of rescaled temperature profiles (for different lattice sizes) for the two models, MC-PIBB, and MC-EBB, respectively. However, note that in the light of the theorem of the previous section, it may be also important to distinguish the cases, of (i) the symmetric interparticle potential with zero pressure ($b = 1, \phi = 0$), and (ii) nonsymmetric case with nonvanishing pressure ($b \neq 1, \phi \neq 0$). We find that for both models (MC-PIBB and MC-EBB), at zero pressure $\phi = 0$, $b = 1$, a non-flat but also non-linear temperature profile is established. However, for nonvanishing pressure, $b = 2.5$, the MC-PIBB model exhibits a vanishing temperature gradient, which is consistent with completely ballistic transport (behavior similar to that found for the integrable models), whereas the MC-EBB model, which is “more ergodic”, even in the case of nonvanishing pressure, establishes a “non-flat” (and non-linear) temperature profile.

From our general theorem of Section IV, we expect that all these momentum conserving bang-bang models should exhibit anomalous conductivity. Using our extensive data on lattices of different sizes, we can confirm this by calculating for each of the models the dependence of the thermal conductivity $\kappa(L)$ on the size L of the system. To account for the observed nonlinearity in the temperature gradients across the system, we have determined “average” temperature gradients ∇T with least squares linear fit of temperature profiles in the range $n = N/4 \dots 3N/4$. Since the temperature profiles are scaling (27), any other choice would just redefine κ by a constant factor (independent of the size L). In Fig.7 we plot $\kappa(L)$ for both lattices (MC-PIBB and MC-EBB) and find significant agreement with the intermediate power-law behavior

$$\kappa(L) \propto L^{0.4}. \quad (28)$$

This result is in agreement with the existing results on non-integrable, momentum-conserving lattices in the literature^{13,20,21,15,33} and provides further support for the conjecture that there may be some form of universality present in these models^{16,15}. Clearly this is an important question worthy of further study.

Our numerical data also permit us to study two other major issues related to the anomalous conductivity and more generally to the applicability of the concepts of statistical mechanics to these systems. First, we can examine the consistency of the above $\kappa(L)$, which we have determined by studying numerically a non-equilibrium steady state, with the behavior of κ as determined by applying the Kubo formula to a large but finite system. To perform this consistency check, we have computed the temporal current-current auto-correlation function, $C(t) \equiv \langle J(t)J \rangle_\beta / L$, and spatio-temporal current-current auto-correlation function, $S(x = m, t) = \langle j_m(t)j_0(0) \rangle_\beta$. The homogeneity in space and time imply that all the averages are invariant under the space and time shifts, $\langle F_m(t) \rangle_\beta = \langle F_0(0) \rangle_\beta$, so the temporal correlation function can be written as the spatial integral of the spatio-temporal one,

$$C(t) = \sum_{m=0}^{N-1} S(m, t). \quad (29)$$

We assume that the tails of the current-current autocorrelation function are mainly governed by the acoustic sound wave propagation which moves ballistically with a group sound velocity $c_s = dx/dt$. This will be clearly revealed for the models studied here by inspecting the full spatio-temporal correlation $S(x, t)$ below.

In order to describe $\kappa(L)$ for a finite system of size L by the conventional Kubo formula, we need to integrate $C(t)$ up to a finite time $t_0 \sim t_L \equiv L/c_s \propto L$, since for the ultimate asymptotic result, the thermodynamic limit $L \rightarrow \infty$ has to be taken prior to the time $t_0 \rightarrow \infty$ limit. Therefore, divergence of $\kappa(L) \propto L^\alpha$ is consistent with a slow power-law decay of current-current autocorrelation function

$$C(t) \propto t^{-(1-\alpha)}. \quad (30)$$

In fig.8 we show temporal correlation functions $C(t)$ for three different models: (a) for MC-PIBB with vanishing pressure ($b = 1$); (b) for MC-EBB with vanishing pressure ($b = 1$);

and (c) for MC-EBB with non-vanishing pressure ($b = 2.5$). Only the case (b) is clearly consistent with the decay $C(t) \propto t^{-0.60}$ whereas for the case (a) (MC-PIBB) the decay of correlations seems to be slightly faster $C(t) \sim t^{-0.78}$ (although the power may approach 0.60 as $N \rightarrow \infty$), and the bumps where the sound-wave collides with its symmetric copy after transversing the half of the system are quite remarkable) and for the case (c) (MC-EBB with non-vanishing pressure), we obtain a finite plateau for $C(t)$, in accordance with our theorem of Section IV.

From these results we conclude that the finite-size relation (30) appears to break down when either (i) the pressure is non-vanishing or (ii) the autonomous dynamics is non-ergodic (pseudo-integrable or even integrable); however, it appears to hold if the pressure is vanishing and the dynamics is ergodic, and of course even more so if dynamics is ergodic and chaotic, as demonstrated by^{20,21}. We believe that our results indicate that EBB models are mixing as well, although it is very difficult to make any precise statements about mixing based on numerical results on decay of time correlations of just one or few observables, like $C(t)$.

Further insight into the nature of the anomalous conductivity comes from studying the full spatio-temporal correlation function $S(x, t)$. In fig.9 we show $S(x, t)$ for the same three cases studied in fig.8. Note the clear tongues of ballistic propagation in $S(x, t)$, from which we can easily compute the sound velocity c_s . For the above three cases the observed sound velocity is: (a) $c_s \approx 1.75$, (b) $c_s \approx 1.67$, (c) $c_s \approx 1.63$. In addition to the ballistic component, we also observe in all three cases a quite pronounced diffusive component of $S(x, t)$ which is the central enhancement (around $x = 0$) inside a band whose width spreads diffusively as $x_{\text{diff}} \propto \sqrt{t}$.

The interpretation of these ballistic modes is clarified by the following simple but instructive observation and numerical experiment. Since there is no on-site potential in our MC-models, they behave macroscopically like a liquid or a gas. When such a system (for a large size $L = N$) is confined between hard walls or between heat reservoirs, we expect it to exhibit standing acoustic waves whose frequencies can be computed directly from the size L and the speed of sound c_s . Since the displacements of the particles must vanish at the

reservoir walls, we also know the appropriate boundary conditions. Thus we should find the same eigen-frequencies as for the acoustic “flute” closed at both ends:

$$\nu_l = \frac{lc_s}{2L}, \quad l = 1, 2, 3 \dots \quad (31)$$

Can we excite the long wavelength acoustic eigenmodes by simple thermal excitation of the reservoirs? We explore this question in Fig.10a by plotting the total displacement q_n of the middle particle (in units of unit mean particle spacings) $n = N/2$ of a MC-EBB chain of size $N = 400$ which is simply put between the heat reservoirs at unit temperature $T_L = T_R = 1$. We observe a nearly periodic signal with a frequency $\nu_1 = 0.0204$, corresponding to a very clear and pronounced dominant excitation of the longest wave-length mode $l = 1$. In the remainder of Fig. 10, we plot the power spectra of the signal, $|\tilde{q}_{N/2}(\omega)|^2$ and $|\tilde{q}_{N/4}(\omega)|^2$, for the four different cases of (see Figs10b-e): (b) a MC-EBB chain with vanishing pressure ($b = 1$), (c) a MC-EBB chain with non-vanishing pressure ($b = 2$), (d) a MC-PIBB chain with vanishing pressure ($b = 1$), and (e) a MC-PIBB chain with non-vanishing pressure ($b = 2$). In all these cases we observe a dominant excitation of the longest wave-length mode $l = 1$ and also clear but weaker excitations of higher eigenmodes ($l > 1$) with integer multiple frequencies $\nu_l = l\nu_1$, whose power is a rapidly decreasing function of l . The basic acoustic frequencies of the four cases are: (b) $2\pi\nu_1 = 0.0128$, (c) $2\pi\nu_1 = 0.0116$, (d) $2\pi\nu_1 = 0.0135$, (e) $2\pi\nu_1 = 0.0116$, from which we can independently calculate the sound velocities: (b) $c_s = 1.63$, (c) $c_s = 1.48$, (d) $c_s = 1.72$, (e) $c_s = 1.48$. The cases (d) and (b) exactly correspond (except for the *different* boundary conditions) to cases (a) and (b) of Figs.8,9, respectively, and indeed the agreement of the sound velocities is very good. Another interesting observation, which we believe is in fact the physical essence of the “proviso” in our theorem of Section IV, is that the power spectra of the “acoustic” signals of models with non-vanishing pressure show a peak (as in Fig.10(c)), or at least relatively large power (as in Fig.10(d)), at zero frequency, whereas models with vanishing pressure have practically vanishing power at zero frequency. In other words: the zero-frequency mode, which is a rigid displacement of all particles (at least in a large local domain, since a global rigid

displacement is prohibited by the boundary conditions), can support energy transport only if the pressure is nonvanishing.

Let us turn now to the second general issue related to the anomalous conductivity and more generally to the applicability of the concepts of statistical mechanics to these systems: namely, is the non-equilibrium state in which we study the heat transport a state of *local thermal equilibrium*? To study the question of the existence of local thermal equilibrium in both PIBB and EBB models in the non-equilibrium stationary state with $T_L = 1, T_R = 2$, we analyze the velocity/momentum distribution across the system and compare it to the ideal Maxwellian,

$$d\mathcal{P}/dp_n \propto \exp\left(-p_n^2/(2m_n T_n)\right).$$

To simplify this comparison, we factor out the local temperature by comparing the normalized higher moments

$$M_{2m}(n) = \frac{\langle p_n^{2m} \rangle}{\langle p_n^2 \rangle^m} \quad (32)$$

with the Gaussian values $M_{2m}^{\text{gauss}} = (2m - 1)!!$.

That we cannot expect to find local thermal equilibrium in a pseudo-integrable (PI) model is shown by the following argument⁴⁶: the momenta of particles in the pseudo-integrable lattice cannot change due to interactions (collisions) with other particles but only due to interactions (collisions) with the reservoirs. Therefore, the velocity distribution at the site n inside the PIBB chain is equal to a linear combination of two Maxwellians with temperatures T_L and T_R , the coefficients being just the probabilities that given velocity has been injected from the left/right. Strictly speaking, for this argument⁴⁶ to be completely justified, we must assume that the diffusion rate of fixed velocities (or “velocitons”) is (at least to some good approximation) independent of the actual value of the velocity. This is indeed the case, for example, if we assume completely uncorrelated random walks for the individual velocities — “velocitons”. But this assumption may be really justified only for strongly chaotic systems, like the Lorentz gas⁴⁶, whereas for PI systems we can consider it just as a qualitative argument that suggests the absence of local thermal equilibrium.

The results of studying the moments of the local velocity distributions in the MC-EBB and MC-PIBB models are shown in Fig.11. The reader should observe very convincing local thermal equilibrium for the MC-EBB model, while for MC-PIBB model we have notable deviations, as anticipated by the qualitative argument presented above.

B. Momentum non-conserving bing-bang models

The two 1D bing-bang models studied in the previous subsection exhibited both translation invariance (momentum conservation) and the absence of metric chaos. Hence the result that they also exhibited anomalous transport is not surprising, given previous results and discussion of Section IV. Nonetheless, our numerics were useful in establishing that the anomalous transport can be viewed as arising from the sound-wave ballistic tongues in spatio-temporal current-current autocorrelation function $S(x, t)$. Further, our simulations also established results—perhaps somewhat surprising, given the complete absence of metric chaos in the bing-bang models—that (i) instead of behaving like integrable systems, the momentum-conserving bing-bang models exhibited the same scaling of $\kappa(L)$ as the generic non-integrable, and even strongly chaotic, models like the FPU lattice; and (2) that in the MC-EBB model, local thermal equilibrium is established in the conducting stationary steady state. Taken together with previous results (for instance, those on the FPU system) that chaos is not sufficient to produce normal conductivity, our new results show that the relationship between metric chaos (positive Lyapunov exponents, positive Kolmogorov-Sinai entropy) and “statistical mechanical” behavior—including normal transport, local thermal equilibrium, it etc.—is perhaps less direct than previously anticipated.

In this subsection, by studying a momentum non-conserving bing-bang model, we further weaken the link between metric chaos and “normal” statistical behavior by establishing numerically that a model in which metric chaos is absent (in the strict sense) nonetheless exhibits normal conductivity: that is, we show by a (numerical) counterexample that chaos is also *not necessary* for normal conductivity.

For simplicity and clarity, we focus on a single momentum non-conserving bing-bang model—the “the less ergodic” pseudo-integrable chain (MNC-PIBB) chain—which is defined by the relations (24) with $b = c = d = 1$, and $m_n = 1$. Physically, this corresponds to a chain of equal mass point particles that collide elastically and are each subject to a hard-wall, confining on-site potential. A typical trajectory of such a model of with nine particles ($N = 9$) is depicted in Fig.12.

When the MNC-PIBB lattice is placed between heat baths, with temperatures $T_1 = 1$ and $T_2 = 2$, a linear temperature profile is established, as shown in Fig. 13, except for the edge particles $n = 1$ and $n = N$, which are in contact with the reservoirs. At the reservoirs, we observe a strong drop in temperature arising from manner in which the particles are coupled to the heat bath: in particular, fast particles are more likely than slow ones to collide again with the reservoirs before they “transmit” their velocity to the rest of the chain. Apart from this effect, which can be essentially avoided by defining “renormalized reservoirs” which include one (or a few) particles near to the reservoir, the transport in the MNC-PIBB is completely *normal*: a linear thermal gradient is established and $\kappa(L)$ is independent of L . And this normal transport behavior occurs despite the fact that the autonomous MNC-PIBB system has no metric chaos and is not even ergodic! We should note, however, that due to pseudointegrability of MNC-PIBB, the local thermal equilibrium is not established in non-equilibrium heat-flow simulation. Instead, the local velocity distribution has been found to be a linear combination of two Maxwellians at different temperatures.

In Fig.14 we show the convergence of $\kappa(L)$ to its L -independent asymptotic value; the figure shows that the convergence has occurred already for lattice sizes $L \approx 100$.

We have conducted three additional numerical experiments to confirm and clarify the normal heat transport in this MNC-PIBB model. First, in fig.16, we show the magnitude of current-current autocorrelation function $|\langle J(t)J \rangle_\beta|/L$ for MNC-PIBB lattices of several different sizes N . We observe a rapid (perhaps exponential) initial decay of the correlations (over three decades) and afterwards a slower, oscillatory decay (so it integrates out and (hopefully) does not produce a divergence in the Kubo formula even if its envelope may not

decay exponentially).

Second, and perhaps more conclusive, is the behavior observed for the spatio-temporal correlation function $S(x, t)$ of the MNC-PIBB model, which we show in Fig.16. In contrast to the behavior observed in the systems with anomalous heat conductivity (see Fig.9), there are *no* ballistic tongues here but instead a clearly diffusive pattern emerges.

Third, we can study the behavior of the imaginary test particles (“velocitons”) that carry constant (fixed) velocity and which can be clearly defined due to pseudo-integrability of the many-body model: When the velocities v_n, v_{n+1} of a pair of particles n and $n + 1$ are exchanged due to the collision, the velociton hops (by definition) from site n to site $n + 1$, or vice versa. Choosing the velociton initially to be at the site $n(t = 0) = n_0$, clearly determines the evolution $n(t)$ is for all times. If the overall motion of the velociton is diffusive, this provides an indication of the diffusive (normal) nature of energy/heat transport in the system. We start by placing a velociton of velocity v_0 somewhere in a large MNC-PIBB lattice (with periodic boundary conditions) and with a canonically thermalized “background”, *i.e.* the momenta of all other particles are distributed according to a Gaussian distribution. We then simulate the dynamics, and ask whether the velociton undergoes a normal diffusive process. If $n(t)$ labels the real particle that carries a velocity v_0 at time t , then we check for the linear growth

$$\langle [n(t) - n(0)]^2 \rangle_\beta = D(v_0)t. \quad (33)$$

In Fig.17 we show $\langle [n(t) - n(0)]^2 \rangle_\beta$ as a function of time, for both an intermediate velociton $v_0 = 1$ (the background temperature is $T = 1$) and a fast velociton $v_0 = 10$. We find normal diffusion in both cases, although the diffusion coefficient $D(v_0)$ is obviously an increasing function of velocity v_0 . In Fig.17 we also study velocitons in a “microcanonically thermalized background”, by which we mean that momenta/velocities of the real particles all have values ± 1 (with equal probability of both signs). Remarkably, even in this case we find normal diffusive processes for both intermediate and fast velocitons.

We believe all these results provide conclusive numerical confirmation of normal diffusive

heat transport in the MNC-PIBB lattice, which is neither ergodic nor chaotic but only pseudo-integrable. This strongly supports the result that *metric chaos is not necessary to have normal transport*.

This result seems quite surprising and indeed paradoxical, but we believe that the resolution of the seeming paradox lies in the complexity of the N -dimensional invariant surfaces in the MNC-PIBB model: namely, these surfaces become more and more dense and uniformly covering of the energy surface as the thermodynamic limit (*i.e.*, as $N \rightarrow \infty$), while dynamics on the invariant surface is probably mixing (decay of time correlations of arbitrary observables), which means that any phase-space distribution function on invariant-surface relaxes into statistical equilibrium. Thus, in the thermodynamic limit $N \rightarrow \infty$, the macroscopic properties of the dynamics of this complex pseudo-integrable system cannot really be distinguished from those of a truly chaotic system. For example, as we saw in the beginning of this section, the sets of modula of momenta of PIBB lattices are preserved under time evolution. Hence the invariant surface can be written as a direct product $(\mathcal{C}_2^N \times \mathcal{S}_N) \times [-1, b]^N$, where $\mathcal{C}_2^N \times \mathcal{S}_N$ is the group of all possible permutations \mathcal{S}_N and sign exchanges \mathcal{C}_2 on a set of initial momenta (p_1, p_2, \dots, p_N) and really represents the discrete momentum part of an invariant surface which consists of $2^N N!$ N -dimensional configurational sheets $[-1, b]^N$. When N is large and if initial condition of on the momenta (\vec{p}) has been chosen with a “canonical measure” for each component p_n (which is the case with probability 1 in thermodynamic limit) then it seems plausible that the dynamics of a single orbit of the PIBB system on its invariant surface can uniformly cover the phase-spaces of small subsystems of size $N' \ll N$ with a canonical measure, just as is the case for a truly ergodic covering system. Obviously, these observations require considerable further mathematical study to be convincing, but we feel that they are well justified as conjectures.

We close this subsection with one final calculation. Since the MNC-PIBB model is equivalent to a (multidimensional) billiard model, we can in this case actually derive the dependence of $\kappa(T)$ on the *temperature* by a simple scaling analysis, either from Kubo formula, or, more directly, from the definition $\kappa = -j/\nabla T$. The heat current is proportional

to the average particle energy, i.e. v^2 (where v is a typical velocity), times the inverse of the typical time in which a particle scatters with its neighbors and transfers energy: this time is roughly a/v , where a is the lattice spacing. Therefore, $j_n \sim v^3$ for billiard-like models. On the other hand, the temperature gradient $\nabla T = T_{n+1} - T_n$ scales as $\sim v^2$, so that $\kappa \propto v$, or

$$\kappa(T) \approx \sqrt{8T} \quad (34)$$

The factor of $\sqrt{8}$ is an approximation obtained from our numerical simulations, but the scaling with temperature is *exact* since it is a simple consequence of scaling dynamics of billiards. The above result is meaningful, of course, only in the close-to-equilibrium situation where typical (average) temperature has a well-defined value of T .

We note that a similar model which exhibits normal heat transport and lacks metric chaos has been recently discussed in Ref.⁴⁷.

VI. SUMMARY AND DISCUSSION

A. Summary

The analyses, analytic and numerical, described in the previous sections, have established several results concerning normal and anomalous heat transport in classical 1D lattices. We have adduced convincing numerical evidence and several theoretical arguments^{22–24} to justify the claim that total momentum conservation is sufficient for anomalous heat transport provided the average pressure is non-vanishing. There seem to be momentum-conserving models that have normal conductivity^{36,37} but these models cannot sustain pressure, consistent with heuristic arguments that in such a case, the Goldstone modes cannot carry energy.

We have also presented numerical results that convincingly support two other claims:

- total momentum conservation is *not* necessary for anomalous heat transport, as shown by the counterexamples of the linear *optical* chain and the nonlinear integrable IK-SG model;

- metric chaos, defined in the usual sense of having a set of non-zero measure in phase space in which there are positive Lyapunov exponents and positive Kolmogorov-Sinai entropy, is *neither* sufficient (as shown by the anomalous behavior of the FPU and diatomic Toda models) *nor* necessary (as shown by the momentum non-conserving pseudo-integrable bing-bang (MNC-PIBB) model) for normal (diffusive) heat transport. Furthermore, even ergodicity is not necessary for normal transport, as shown by our studies of MNC-PIBB models. From our analysis it follows that perhaps even multidimensional pseudo-integrability is sufficient, provided the topology of invariant surfaces becomes sufficiently complex in thermodynamic limit.

As is often the case, these results raise at least as many questions as they answer, so we phrase the remainder of our discussion in terms of several questions.

B. What is needed for normal transport?

Our results establish that the full set of conditions needed to guarantee normal transport in 1D models is more subtle and less “clean” than previously believed⁷. We know that normal transport for all temperatures requires

- Confining on-site potential¹⁶, or vanishing pressure with effective means of scattering of long-wavelength Fourier modes^{36,37}.
- Anharmonicity in either on-site or interparticle potential: this is necessary to exclude the pathological linear systems;
- Absence of integrability: integrable models (which can be hard to identify *a priori* !) must also clearly be excluded;
- An *effective* means of achieving dynamical mixing in the thermodynamic limit, although this means can be subtle and perhaps more topological than dynamical, as the results for the pseudo-integrable bing-bang model imply.

C. What about dimensionality?

Why do we expect our arguments to be limited to one dimension? We could develop a formal analysis of the vector current in two (or higher) dimensions, but intuition gained from thinking of coupled chains is key. The neighboring chains provide an environment for a given chain that leads to an effective on-site potential: this was original motivation for Frenkel-Kontorova model. Based on hydrodynamic or mode-coupling arguments⁴⁸, one would expect $\ln L$ divergence of κ in 2D and normal conductivity $\kappa \propto L^0$ in 3D. The expected logarithmic divergence in 2D has been confirmed in recent numerical experiments³³. Nonetheless, further studies in explicit lattice models in higher dimensions will certainly provide additional insights.

D. What are other open issues?

- An outstanding problem in mathematical physics is a rigorous proof of what is really necessary and sufficient for normal conductivity.
- Compelling calculations of the exponents for anomalous transport. Is $\kappa \sim L^{1.0}$ the fastest increase possible? Is α universal, or are there universality classes? Numerical data suggest $\alpha \sim 0.4$ is widespread, but other arguments suggest $\alpha = 1/3$. Is there a systematic “universality theory” that can predict α , as was done for the Feigenbaum constants in the period doubling transition to chaos?
- What happens when systems are driven *non*-linearly away from equilibrium? What are the corrections to Kubo formula?

Knowledgeable colleagues, many of whom have contributed to this Focus Issue of *Chaos* on the Fermi-Pasta-Ulam problem, can doubtless add still further questions. There is still much to be found in exploring the rich trove of physically relevant and mathematically challenging problems that has been uncovered in seeking to explain FPU’s remarkable “little discovery”¹.

ACKNOWLEDGMENTS

We thank J. L. Lebowitz, G. Casati, Bambi Hu, Peter Wolynes, and C. Mejia-Monasterio for fruitful discussions. We are especially grateful to the Center for Nonlinear Studies (CNLS) at Los Alamos National Laboratory, whose hospitality and support enabled us to begin this collaboration. DKC acknowledges the support provided by the Ulam Scholar position at the CNLS and by the visitors program at MIDIT at Danish Technical University. TP thanks the Centro Internacional de Ciencias in Cuernavaca, Mexico, for hospitality and the Ministry of Science and Technology of Slovenia for financial support.

REFERENCES

- ¹ E. Fermi, J. Pasta, S. Ulam, “Studies of Nonlinear Problems. I” Los Alamos report LA-1940 (1955); reprinted in E. Fermi. *Collected Papers*, Vol II (Univ. of Chicago Press. Chicago, 1965), p. 978.
- ² Morikazu Toda, “Studies of a Non-Linear Lattice,” *Phys. Rep.* **18**, 1-124 (1975).
- ³ W. M. Visscher, “Computer Studies of Transport Properties of Simple Models of Solids” *Meth. Comput. Phys.* **15**, 371-408 (1976).
- ⁴ R. A. MacDonald and D. H. Tsai, “Molecular Dynamical Calculations of Energy Transport in Crystalline Solids,” *Phys. Rep.* **46**, 1-41 (1978).
- ⁵ E. Atlee Jackson, “Nonlinearity and Irreversibility in Lattice Dynamics,” *Rocky Mountain J. of Math.* **8** 127-196 (1978).
- ⁶ Morikazu Toda, “Solitons and Heat Conduction,” *Phys. Scr.* **20**, 424-430 (1979).
- ⁷ Joseph Ford, “The Fermi-Pasta-Ulam Problem: Paradox turns Discovery,” *Phys. Rep.* **213**, 271-310 (1992).
- ⁸ R. Livi, A. Politi and S. Lepri, “Thermal conduction in classical low-dimensional lattices”, *Physics Reports* **377**, 1-80 (2003).
- ⁹ *Transport Phenomena*, Springer Lecture Notes in Physics Vol. 31, G. Kirczenow and J. Marro, eds (Springer Verlag, 1974) remains a classic reference for discussion of all these issues.
- ¹⁰ W. M. Visscher, “Transport processes in solids and linear response theory”, *Phys. Rev. A* **10**, 2461 (1974).
- ¹¹ Z. Rieder, J. L. Lebowitz, and E. Lieb, “Properties of a Harmonic Crystal in a Stationary Nonequilibrium State,” *J. Math. Phys.* **8** 1073-1078 (1967).
- ¹² Hideo Kaburaki and Masahiko Machida, “Thermal conductivity in one-dimensional lat-

- tices of the Fermi-Pasta-Ulam type,” *Phys. Lett. A* **181**, 85-80 (1993).
- ¹³ Stefano Lepri, Roberto Livi, and Antonio Politi, “Heat Conduction in Chains of Nonlinear Oscillators,” *Phys. Rev. Lett.* **78**, 1896-1899 (1997).
- ¹⁴ E. A. Jackson and A. D. Mirlis, “Thermal conductivity on one- and two-dimensional lattices,” *J. Phys. Cond. Matt.* **1**, 1223 (1989).
- ¹⁵ Takahiro Hatano, “Heat conduction in the diatomic Toda lattice revisited,” *Phys. Rev. E* **59**, R1-R4 (1999).
- ¹⁶ Bambi Hu, Baowen Li, and Hong Zhao, “Heat conduction in one-dimensional chains,” *Phys. Rev. E* **57** 2992-2995 (1998).
- ¹⁷ G. Casati, J. Ford, F. Vivaldi, and W. H. Visscher, “One-dimensional classical many-body system having normal thermal conductivity”, *Phys. Rev. Lett.* **52**, 1861-1864 (1984).
- ¹⁸ Tomaž Prosen and Marko Robnik, “Energy Transport and detailed verification of Fourier heat law in a chain of colliding harmonic oscillators,” *J. Phys A: Math. Gen.* **25**, 3449-3472 (1992).
- ¹⁹ Robert S. Northcote and Renfrey B. Potts, “Energy Sharing and Equilibrium in Nonlinear Systems,” *J. Math. Phys.* **5**, 383-398 (1964).
- ²⁰ Stefano Lepri, Roberto Livi, and Antonio Politi, “Energy Transport in anharmonic lattices close to and far from equilibrium,” *Physica D* **119**, 140-147 (1998).
- ²¹ Stefano Lepri, Roberto Livi, and Antonio Politi, “On the anomalous thermal conductivity of one-dimensional lattices,” *Europhys. Lett.* **43** 271-276 (1998).
- ²² Tomaž Prosen and David K. Campbell, *Phys. Rev. Lett.* **84** 2857-2860 (2000).
- ²³ O. Narayan and S. Ramaswamy, “Anomalous heat conduction in one-dimensional momentum-conserving systems”, *Phys. Rev. Lett.* **89**, 200601 (2002).

- ²⁴ G. Casati and T. Prosen, “Anomalous heat conduction in a one-dimensional ideal gas”, Phys. Rev. E **67**, 015203(R) (2003).
- ²⁵ F. Bonetto, J. L. Lebowitz and L. Rey-Bellet, “Fourier law: a challenge to theorists”, preprint *arXiv:math-ph/0002052*.
- ²⁶ A. Dhar, “Heat conduction in the disordered harmonic chain revisited”, Phys. Rev. Lett. **86**, 5882-5885 (2001).
- ²⁷ Morikazu Toda, “Vibration of a Chain with Nonlinear Interaction,” *J. Phys. Soc. Jpn.* **22**, 431-436 (1967).
- ²⁸ F. Mokross and H. Büttner, J. Phys. C **16**, 4539 (1983).
- ²⁹ Brad Lee Holian, Hermann Flaschka, and David W. McLaughlin, “Shock Waves in the Toda lattice: Analysis,” *Phys. Rev. A* **24**, 2595-2623 (1981); B. L. Holian and G. K. Straub, “Molecular dynamics of shock waves in one-dimensional chains,” *Phys. Rev. B* **18**, 1593-1608 (1978); G. K. Straub, B. L. Holian, and R. G. Petschek, “Molecular dynamics of shock waves in one-dimensional chains. II: Thermalization,” *Phys. Rev. B* **19**, 4049-4055 (1979).
- ³⁰ A. G. Izergin and V. E. Korepin, “A lattice sine-Gordon model,” *Vestn. Leningr. Gos. Univ., Fiz. Khim.* **22**, 88 (1981).
- ³¹ V. O. Tarasov, “A Classical Version of the lattice sine-Gordon model,” Plenum translation 1986 (need to get exact reference) 2011-2022 (1986) : [*Zap. Nauk. Sem. Lenin. Otdel. Mat. Inst. im. V. A. Steklova* **120** 173-187 (1982).]
- ³² W. H. Press, B. P. Flannery, S. A. Teukolsky and W. T. Vetterling, “Numerical Recipes”, (Cambridge University Press, 1988).
- ³³ A. Lippi, Diploma Thesis, University of Florence 1998.
- ³⁴ P. Grassberger, W. Nadler and L. Yang, “Heat conduction and entropy production in a

- one-dimensional hard-particle gas”, *Phys. Rev. Lett.* **89**, 180601 (2002).
- ³⁵ A. Dhar, “Heat conduction in a one-dimensional gas of elastically colliding particles of unequal masses”, *Phys. Rev. Lett.* **86**, 3354-3357 (2001).
- ³⁶ C. Giardinà, R. Livi, A. Politi and M. Vassalli, “Finite Thermal Conductivity in 1D Lattices”, *Phys. Rev. Lett.* **84**, 2144-2147 (2000).
- ³⁷ O. V. Gendelman and A. V. Savin, “Normal Heat Conductivity of the One-Dimensional Lattice with Periodic Potential of Nearest-Neighbor Interaction”, *Phys. Rev. Lett.* **84**, 2381-2385 (2000).
- ³⁸ J. M. Deutsch and O. Narayan, “One dimensional heat conductivity exponent from a random collision model”, *Phys. Rev. E* **68**, 010201(R) (2003).
- ³⁹ J. M. Deutsch and O. Narayan, “Correlations and scaling in one-dimensional heat conduction”, *Phys. Rev. E* **68**, 041203 (2003).
- ⁴⁰ L. Yang and P. Grassberger, “Are there really phase transitions in 1-d heat conduction models?”, preprint [arXiv:cond-mat/0306173](https://arxiv.org/abs/cond-mat/0306173).
- ⁴¹ R. Kubo, M. Toda and N. Hashitsume, “Statistical Physics II”, Springer Series in Solid State Sciences, vol. 31, (Springer-Verlag, 1991).
- ⁴² E. Gutkin, “Billiards in polygons: Survey of recent results”, *J. Stat. Phys.* **83**, 7-26 (1996).
- ⁴³ G. Casati and T. Prosen “Mixing property of triangular billiards”, *Phys. Rev. Lett.* **83**, 4729-4732 (1999).
- ⁴⁴ G. Casati and T. Prosen, “Triangle map: a model of quantum chaos”, *Phys. Rev. Lett.* **85**, 4261-4264 (2000).
- ⁴⁵ G. Casati, “Energy Transport and the Fourier Heat Law in Classical Systems” *Foundations of Physics* **16**, 51-61 (1985).

- ⁴⁶ The same argument has been used recently for a one-particle problem, namely the Lorentz gas. See A. Dhar and D. Dhar, “Absence of local thermal equilibrium in two models of heat conduction”, *Phys. Rev. Lett.* **82**, 480-483 (1999).
- ⁴⁷ B. Li, G. Casati, J. Wang and T. Prosen, “Fourier law in the alternate-mass hard-core potential chain”, *Phys. Rev. Lett.* **92**, 254301 (2004).
- ⁴⁸ see e.g. Peter G. Wolynes, “Bounds for convective contributions to transport coefficients,” *Phys Rev A* 11, 1700-1705 (1975); Peter G. Wolynes, “Hydrodynamic boundary conditions and mode-mode coupling theory,” *Phys Rev A* 13, 1235-1250 (1976); Y. Pomeau and P. Resibois, “Time dependent correlation functions and mode-mode coupling theories”, *Physics Reports* **19**, 63-139 (1975); M. H. Ernst, “Mode-coupling theory and tails in CA fluids”, *Physica D* **47**, 198-211 (1991).

FIGURES




FIG. 1. Temperature profile for the linear *optical* chain with the parameter $\tilde{\nu} = 0.25$ and with reservoir temperatures $T_L = 1.3, T_R = 0.7$. The size of the lattice here is $N = 40$. As discussed in the text, this profile is the same as for the linear acoustic chain discussed in Ref. ⁽¹¹⁾




FIG. 2. The width, $\sigma(t) \equiv \sqrt{(\sum_n n^2 h_n)/(\sum_n h_n)}$, of a pulse spreading from an initial single-site disturbance as a function of time in the IKSG model for two different values of parameter $b = \tan \delta$. The initial excitation is delta-like with $\phi_0^+ = 2, \phi_0^- = 3, \phi_n^\pm = 0, n \neq 0$. A finite lattice of length $N = 800$ with periodic boundary conditions has been used in the simulation. The line with slope 1 is used to guide the eye: the figure provides clear evidence of ballistic transport, $\sigma(t) \sim t$.




FIG. 3. Typical trajectories of the individual particles in a MC-PIBB model with constant unit masses and size $L = 9$. The ordinate shows particle positions ($x_n(t)$) where for clarity the odd n particle trajectories are plotted with full lines while the even n trajectories are dashed. The walls at $x = 0$ and $x = 10$ act as thermal baths at temperature $T = 1$.




FIG. 4. Typical trajectory of a MC-EBB model with mass ratio $m_2/m_1 = 0.382$ and size $L = 9$. As in Fig.3, the ordinate shws particle positions ($x_n(t)$) where for clarity the odd n particle trajectories are plotted with full lines while the even n trajectories are dashed. The walls at $x = 0$ and $x = 10$ act as thermal baths at temperature $T = 1$. .




FIG. 5. Scaling of temperature profiles (temperature vs. x/N for different system sizes N) in a MC-PIBB model with thermal baths at temperatures $T_L = 0.5, T_R = 1.0$. The triple of steep curves refers to a system with symmetric interparticle potential $b = 1.0$ (so that the pressure vanishes), while the triple of horizontal curves (at $T = \sqrt{T_L T_R}$, so zero temperature gradient over the bulk of the chain) refers to $b = 2.5$ (and a non-vanishing pressure).




FIG. 6. Scaling of temperature profiles (temperature vs. x/N for different system sizes N) in a MC-EBB model with mass ratio $m_2/m_1 = 0.382$. All other parameters are the same as in Fig.5. Note that here a nonvanishing temperature gradient is established even in the non-symmetric case $b = 2.5$ with nonvanishing pressure.






FIG. 7. The finite-size thermal conductivity $\kappa(L)$ vs. size L for the MC-EBB ($m_2/m_1 = 0.382, b = 1$) and MC-PIBB ($m_2/m_1 = 1, b = 1$) models.










FIG. 8. The current-current temporal autocorrelation function $\langle J(t)J \rangle_\beta / L$ (at temperature $T = 1$) for: (a) the pseudointegrable ($m_2/m_1 = 1$) momentum-conserving bing-bang model with periodic boundary conditions and vanishing pressure, $b = 1.0$; (b) the ergodic momentum-conserving bing-bang lattice ($m_2/m_1 = 0.382$) with vanishing pressure, $b = 1.0$; and (c) the ergodic momentum-conserving lattice with non-vanishing pressure $b = 2.5$. In each case, three different system sizes are shown. Note that the dotted line in (c) indicates the lower bound from the theorem from section IV.












FIG. 9. The spatiotemporal correlation function $S(m, t) = \langle j_m(t) j_0(0) \rangle_\beta$ for the same three cases as in Fig.8. Note the apparent ballistic propagation. Regions between contours of equidistantly spaced base- e logarithm $\ln S(x = m, t)$ are shaded with 20 different and uniformly increasing levels of greyiness in the range $[-10, 0]$.












FIG. 10. Thermal excitation of acoustic modes: (a) the time dependence of the displacement $q_n(t)$ of the particle in the middle $n = L/2$ of the diatomic ($m_2/m_1 = 0.382, b = 1.0$) bing-bang lattice of size $L = 400$ between thermal reservoirs in equilibrium ($T_L = T_R = 1$); (b) the power-spectrum of the same signal ($n = L/2$, full curve) and for the displacements at the one-quarter of the chain ($n = L/4$, dashed curve); dotted vertical lines denote integer multiples of the basic acoustic frequency; (c) the same plot as in (b) but for the MC-EBB with $b = 2$ (nonvanishing pressure); (d) the same plot as in (b) but for the MC-PIBB with $b = 1$ (vanishing pressure); and (e) the same plot as in (b) but for the MC-PIBB with $b = 2$ (non-vanishing pressure).




FIG. 11. Moments of the velocity distributions $M_{2n}(m) = \langle p_m^{2n} \rangle_\beta / \langle p_m^2 \rangle_\beta^n$ for $n = 4, 6$, and 8 for two different momentum conserving bing-bang lattices of size $L = 3200$ with $T_L = 0.5, T_R = 1.0$. The results for the EBB model (with $m_2/m_1 = 0.382$, plotted as solid lines) are essentially indistinguishable from those of the Gaussian model ($M_n = (2n - 1)!!$), plotted as dotted lines), showing that local thermal equilibrium is established in the MC-EBB model, whereas the results for the MC-PIBB model (dashed lines) deviate substantially from the Gaussian model, showing that local thermal equilibrium is *not* established, as anticipated by the heuristic argument presented in the text.




FIG. 12. A typical trajectory of MNC-PIBB chain ($L = 9$). As in Fig.3, the ordinate shows particle positions ($x_n(t)$) where for clarity the odd n particle trajectories are plotted with full lines while the even n trajectories are dashed. The walls at $x = 0$ and $x = 10$ act as thermal baths at temperature $T = 1$.




FIG. 13. The scaling of *linear* temperature profiles, T vs x/L , for MNC-PIBB chain for different sizes L . The dashed-dotted straight line is drawn to guide the eye. The temperatures of stochastic heat baths are $T_L = 1$ and $T_R = 2$.




FIG. 14. Convergence of the thermal conductivity $\kappa(L)$ to its L -independent asymptotic value for the MNC-PIBB model. In the simulations, $T_L = 1$, and $T_R = 2$.




FIG. 15. The magnitude of the current autocorrelation function $|\langle J(t)J \rangle|/L$ (at temperature $T = 1$) for the MNC-PIBB model with periodic boundary conditions for different system sizes.




FIG. 16. The spatiotemporal correlation function $S(m, t) = \langle j_m(t) j_0(0) \rangle_\beta$ for the same parameters as in previous Fig.15 and using the same graphical presentation as in Fig.9. Note the apparent diffusive propagation. The superimosed full curve is $t = 0.09x^2$.




FIG. 17. Diffusion of ‘velocitons’ in MNC-PIBB lattice. We show root-mean-square displacements vs. time for velocitons with $v = 1$ and $v = 10$, in either canonically thermalized background with temperature $T = 1$ or in ‘microcanonically thermalized’ background (all background particles have equal unit velocities but random signs and initial positions). Full lines give the slopes of best linear fits which support the existence of normal diffusion.